

NOTE

A Bijective Proof of Lassalle's Partition Identity

Jiang Zeng

*Institut Girard Désargues, Université Claude Bernard Lyon 1,
43, boulevard du 11 Novembre 1918, 69622 Villeurbanne Cedex, France*
E-mail: zeng@desargues.univ-lyon1.fr

Communicated by the Managing Editors

Received June 7, 1999; accepted June 17, 1999

[View metadata, citation and similar papers at core.ac.uk](#)

We give a bijective proof of the following form of a conjecture in [2, 3]. The special $s = 1$ case was proved in the same vein by Simion [2]. For other proofs of this conjecture see [4]. Our notations are the same as in [2, 3]. For an integer partition μ let the length $l(\mu)$ be the number of the parts of μ , $m_i(\mu)$ the number of parts equal to i , $z_\mu = \prod_{i \geq 1} i^{m_i(\mu)} m_i(\mu)!$ and $\langle \mu_r \rangle$ the number of ways to choose r different cells from the diagram of the partition μ taking at least one cell from each row. For any positive integer n let $(a)_n = a(a+1) \cdots (a+n-1)$ and $[n] = \{1, 2, \dots, n\}$.

THEOREM 1. *We have*

$$\sum_{|\mu|=n} \frac{n!}{z_\mu} \langle \mu_r \rangle X^{l(\mu)-1} \sum_{i=1}^{l(\mu)} \mu_i \binom{\mu_i + s - 1}{s - 1} \\ = \binom{n}{r} (s+r)_{n-r} [(X+s)_r - (X)_r]. \quad (1)$$

Recall that the functional graph of an injection $f: [n] \rightarrow [n+r]$ consists of disjoint cycles and paths. Each path is of form $a_0 \rightarrow a_1 \rightarrow \cdots \rightarrow a_l$, where $f(a_i) = a_{i+1}$ for $0 \leq j < l$, with $f^{-1}(a_0)$ empty, and $a_l \in [n+r] \setminus [n]$. Each cycle is of the form $a_0 \rightarrow a_1 \rightarrow \cdots \rightarrow a_l$, where $f(a_i) = a_{i+1}$ for $0 \leq j \leq l$, with $a_{l+1} = a_0$. Let $\text{cyc } f$ be the number of cycles in f . Then the corresponding generating function reads as follows:

$$\sum_{f: [n] \rightarrow [n+s]} X^{\text{cyc } f} = (X+s)_n \quad (2)$$

where f is an injection. The above identity seems to be first published by Foata and Strehl [1] and can be readily verified by induction on n as in the well-known case of permutations, which corresponds to $s=0$.

We shall prove that the two sides of (1) are the generating function of pairs (f, R) with respect to the number of cycles of f , where f is an injection from $[n]$ to $[n+s]$ and R is a r -subset of $[n]$ such that each cycle of f has at least one element in R .

Indeed, to construct such a pair (f, R) we can first choose r elements of $[n]$ to form a r -subset R , there are $\binom{n}{r}$ ways to do so; and construct an injection $g: [r] \rightarrow [r+s]$ such that its restriction on $[r]$ is not a permutation of $[r]$, the generating function of such injections is then $(X+s)_r - (X)_r$; finally insert successively the $n-r$ elements of $[n] \setminus R$ into the above injection g to obtain the injection $f: [n] \rightarrow [n+s]$, there are $(s+r)_{n-r}$ ways to insert the $n-r$ elements into any above injection g . This gives the right-hand side of (1).

On the other hand, we can construct such a pair (f, R) as follows: construct first a permutation σ on $[n]$, for any partition μ of n there are $n!/z_\mu$ such permutations of type μ ; choose r elements of $[n]$ to form the subset R such that each cycle of σ has at least one element in R , there are $\langle \mu_r \rangle$ ways to do so; choose any cycle $(a, \sigma(a), \dots, \sigma^{\mu_i-1}(a))$ of σ with length μ_i to form s paths:

$$\begin{aligned} a &\rightarrow \sigma(a) \rightarrow \dots \sigma^{l_1-1}(a) \rightarrow n+1, \\ \sigma^{l_1}(a) &\rightarrow \sigma^{l_1+1}(a) \rightarrow \dots \sigma^{l_2-1}(a) \rightarrow n+2, \\ &\dots \rightarrow \dots \\ \sigma^{l_{s-1}}(a) &\rightarrow \sigma^{l_{s-1}+1}(a) \rightarrow \dots \sigma^{l_s-1}(a) \rightarrow n+s, \end{aligned}$$

where $l_1 + l_2 + \dots + l_s = \mu_i$ and by convention the corresponding path is empty if $l_i = 0$. Obviously there are $\mu_i \binom{\mu_i + s - 1}{s-1}$ ways to form such s paths starting from any cycle of length μ_i . Finally we complete the construction of $f: [n] \rightarrow [n+s]$ by using the s paths above and the remaining $l(\mu) - 1$ cycles of σ . This gives the left-hand side of (1).

The theorem follows then from the two above countings.

REFERENCES

1. D. Foata and V. Strehl, Combinatorics of Laguerre polynomials, in "Enumeration and Design, Waterloo Jubilee Conference," pp. 123–140, Academic Press, New York, 1984.
2. M. Lassalle, Quelques conjectures combinatoires relatives à la formule classique de Chu–Vandermonde, *Adv. in Appl. Math.* **21** (1998), 457–472.
3. M. Lassalle, Une conjecture en théorie des partitions, <http://xxx.lanl.gov/math.CO/9901040>, 1999.
4. M. Lassalle, Une identité en théorie des partitions, *J. Combin. Theory Ser. A* **89** (2000), 270–288, doi:10.1006/jcta.1999.3010.